

# Attribution of Seasonal Climate Anomalies August-September-October 2024

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


# Summary of Observed Conditions and Outlooks

- In ASO2024, weakly cold SST anomalies were observed in the central-eastern equatorial Pacific and extended southward along the coastal regions of South America. For most of the other ocean basins - North Pacific, central southern Pacific, tropical and north Atlantic, and Indian Ocean – warm SST anomalies persisted (slide 4). CFSv2 forecast the large-scale structure of observed SST anomalies but had a cold prediction bias over tropical and northern Pacific and North Atlantic in general (Slide 10). Tropical Atlantic has remained abnormally warm for over a year.
- The AMIP simulation and the initialized CFSv2 forecast, as well as other NMME forecasts had positive anomalous rainfall over the Maritime Continent extending into the southwestern Pacific and dry conditions over the central and eastern equatorial Pacific, a pattern similar to [the canonical La Nina response](#). The forecast pattern, however, differed from the observations and a less expansive coverage for negative anomalies (Slide 11, 37-39).
- Consistent with the continued warming of SSTs globally, the tendency for above normal 200-mb heights and above normal land surface temperature continued to dominate across the globe both in observations and model predictions (slide 10, 12, 13).
- The initialized CFSv2 forecasts generally well predicted the tendency for above normal 200-mb height and land surface temperature over North America (slide 15, 16).
- The initialized CFSv2 forecasts captured most of the observed dry precipitation anomalies across the central US extending into Mexico, as well as the wet precipitation anomalies in the US Southeast (slide 14).
- October 2024 monthly forecast skill for North America's T2m and precipitation show improvement from the shortest leads compared to the longer leads (slide 34-35).


# Observed Seasonal Anomalies

## Global and North America


# Observed Anomaly AS02024



# Observed Anomaly AS02024






# Observed Anomaly AS02024




# CPC Seasonal Outlooks and NMME Forecasts

CPC


Temp nonEC  
HSS=89



Prec nonEC  
HSS=38




NMME prob fcst TMP2m IC=202407 for lead 1 2024 ASO



NMME

NMME prob fcst Prate IC=202407 for lead 1 2024 ASO



For the rationale behind CPC outlooks see [https://www.cpc.ncep.noaa.gov/products/archives/long\\_lead/PMD/2024/202407\\_PMD90D](https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2024/202407_PMD90D)


## Model Simulated/Forecast Ensemble Mean Anomalies




# Model Simulated/Forecast Ensemble Average Anomalies

- **AMIP simulations** forced with observed sea surface temperatures (100 members ensemble)
- CFSv2 real time operational forecasts
  - **Seasonal forecast**: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
  - **Reconstructed forecast**: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC). “w/o mean” is AC with area mean removed.


# ASO2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




# ASO2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




# AS02024 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)




# AS02024 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




# AS02024 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



# AS02024 Observed & Model Simulated/Forecast Ensemble Average Anomalies z200(m)



# AS02024 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)






## Model Simulated/Forecast Anomalies: Individual Runs

## Model Simulated/Forecast Anomalies: Individual Runs


- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).

# ASO2024 Anomaly Correlation for Individual AMIP Simulation with Observation -- z200(20N-90N)




# Observed & AMIP Ensemble Mean Anomalies


## ASO2024 z200(m) 100 runs/worst 4 runs/best 4 runs




# ASO2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N-90N)




Observed & CFSv2 Forecast Ensemble Average Anomalies  
ASO2024 z200(m) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast



Observed & CFSv2 Forecast Ensemble Average Anomalies  
AS02024 z200(m) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast




# ASO2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S-30N)







Observed & CFSv2 Forecast Ensemble Average Anomalies  
ASO2024 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies  
AS02024 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast




# ASO2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S-30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies  
ASO2024 T2m(K) 40 runs/worst 4 runs/best 4 runs  
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies  
ASO2024 T2m(K) 40 runs/worst 4 runs/best 4 runs  
Reconstructed Forecast



# z200(m) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) AS02024 z200(m) eddy & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2<sup>nd</sup> row: last 10 days of the prior month.
- 3<sup>rd</sup> row: 11<sup>th</sup> - 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 1<sup>st</sup> - 10<sup>th</sup> of the prior month.

# T2m(k) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) ASO2024 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2<sup>nd</sup> row: last 10 days of the prior month.
- 3<sup>rd</sup> row: 11<sup>th</sup> - 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 1<sup>st</sup> - 10<sup>th</sup> of the prior month.

# Prec(mm/day) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) ASO2024 Prec(mm/day) & Obs



Top row: Observed anomaly.


CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2<sup>nd</sup> row: last 10 days of the prior month.
- 3<sup>rd</sup> row: 11<sup>th</sup> - 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 1<sup>st</sup> - 10<sup>th</sup> of the prior month.



# z200(m) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst ASO2024 z200(m) eddy & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2<sup>nd</sup> row: last 3 days of the prior month.
- 3<sup>rd</sup> row: 18<sup>th</sup> – 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 7<sup>th</sup> – 10<sup>th</sup> of the prior month.

# T2m(k) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst ASO2024 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2<sup>nd</sup> row: last 3 days of the prior month.
- 3<sup>rd</sup> row: 18<sup>th</sup> – 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 7<sup>th</sup> – 10<sup>th</sup> of the prior month.

# Prec(/mm/day) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst ASO2024 Prec(mm/day) & Obs



Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2<sup>nd</sup> row: last 3 days of the prior month.
- 3<sup>rd</sup> row: 18<sup>th</sup> – 20<sup>th</sup> of the prior month.
- 4<sup>th</sup> row: 7<sup>th</sup> – 10<sup>th</sup> of the prior month.



## Seasonal Forecasts from Multi-Model Ensemble Systems

- WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME).  
<https://www.wmolc.org/>
- Copernicus Climate Change Service (C3S) Multi-model seasonal forecasts.  
[https://climate.copernicus.eu/charts/c3s\\_seasonal/](https://climate.copernicus.eu/charts/c3s_seasonal/)
- North American Multi-Model Ensemble (NMME) seasonal forecasts.  
<https://www.cpc.ncep.noaa.gov/products/NMME/>



# LC-LRFMM Seasonal Forecasts

(<https://www.wmolc.org/>)

## Ensemble means



## Probabilities



# C3S Seasonal Forecast

([https://climate.copernicus.eu/charts/c3s\\_seasonal/](https://climate.copernicus.eu/charts/c3s_seasonal/))

C3S multi-system seasonal forecast

Mean forecast SST anomaly


Nominal forecast start: 01/07/24

Variance-standardized mean

ECMWF/Met Office/Météo-France/CMCC/NCEP/JMA

ASO 2024

Legend for SST anomaly: <math>< -2.0^{\circ}\text{C}</math>, <math>-2.0 \dots -1.0</math>, <math>-1.0 \dots -0.5</math>, <math>-0.5 \dots -0.2</math>, <math>-0.2 \dots 0.2</math>, <math>0.2 \dots 0.5</math>, <math>0.5 \dots 1.0</math>, <math>1.0 \dots 2.0</math>, <math>> 2.0^{\circ}\text{C}</math>



C3S multi-system seasonal forecast

Mean precipitation anomaly


Nominal forecast start: 01/07/24

Variance-standardized mean

ECMWF/Met Office/Météo-France/CMCC/NCEP/JMA



ASO 2024

Legend for precipitation anomaly: <math>< -200\text{mm}</math>, <math>-200 \dots -100</math>, <math>-100 \dots -50</math>, <math>-50 \dots -10</math>, <math>-10 \dots 10</math>, <math>10 \dots 50</math>, <math>50 \dots 100</math>, <math>100 \dots 200</math>, <math>> 200\text{mm}</math>





# North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)



# 200mb Height from Linear Model

ASO2024 200mb Eddy HGT(m)  
OBS vs. Linear Model Response to Tropical Heating  
Heating is converted from Prate in 15S-15N




Pattern COR: global=0.15, tropics(30S-30N)=0.28



# Seasonal Forecasts from the Constructed Analog Model


CA HGT200 Prd for ASO2024, ICs through Oct2024(m), Lead -3



Michael Goss NOAA/NWS/NCEP/CPC

Base Period 1931-2020


CA T2m Prd for ASO2024, ICs through Oct2024(K), Lead -3



Michael Goss NOAA/NWS/NCEP/CPC

Base Period 1931-2020

CA Prec Prd for ASO2024, ICs through Oct2024(mm/day), Lead -3



Michael Goss NOAA/NWS/NCEP/CPC

Base Period 1931-2020

## Background & Methodology

# Attribution of Seasonal Climate Anomalies

- Goal
  - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
  - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

# Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
  - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
  - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

## Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

# Data

- Observations
  - SST: OI version 2 analysis (Reynolds et al., 2007)
  - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
  - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
  - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
  - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
  - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation based on GFS\_FV3 (provided by Dr. Tao Zhang/CPC)
  - 100 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model.
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model.