

Attribution of Seasonal Climate Anomalies

September-October-November 2024

(<https://www.cpc.ncep.noaa.gov/products/people/mchen/AttributionAnalysis/>)


Summary of Observed Conditions and Outlooks

- In SON2024, cold SST anomalies in the central-eastern and far eastern equatorial Pacific remained marginal. Meanwhile, warm SST anomalies persisted across most other ocean basins, including the North Pacific, central and southern Pacific, tropical and North Atlantic, and Indian Ocean (slide 4). The CFSv2 model captured the large-scale structure of observed SST anomalies but show a tendency toward cold prediction bias over the tropical and northern Pacific, North Atlantic, and Indian Ocean (slide 10). The tropical Atlantic continued to stay anomalously warm for over a year.
- The AMIP simulation and the initialized CFSv2 forecast, as well as other MME forecasts had positive anomalous rainfall over the Maritime Continent extending into the southwestern Pacific and dry conditions over the central and eastern equatorial Pacific, a pattern similar to [the canonical La Niña response](#). The forecast pattern, however, differed from the observations and had a much smaller coverage for negative anomalies (Slide 11, 37-39).
- The persistence of above-normal anomalies in 200-mb heights and land surface temperatures reflects the atmospheric response to continued global SST warming and was consistent with warming over the global oceans acting as the forcing. The model successfully captured the observed warming trend and the large-scale structure of 200-mb height eddy component anomalies (slides 10, 12, 13).
- The initialized CFSv2 forecasts captured the overall tendency for above normal 200-mb height and land surface temperature over North America. However, discrepancies remain, as the centers of eddy anomalies show some spatial shifts compared to observations (slides 15, 16).
- The initialized CFSv2 forecasts captured most of the observed dry precipitation anomalies across the northern central US and Mexico, and wet conditions in the southeastern US (slide 14).
- November 2024 monthly forecast skill for North America's 200-mb height, T2m, and precipitation did not show improvement from the shortest leads compared to the longer leads (slide 34-35).


Observed Seasonal Anomalies

Global and North America


Observed Anomaly SON2024



Observed Anomaly SON2024






Observed Anomaly SON2024





CPC Seasonal Outlooks and NMME Forecasts


CPC



Prec nonEC
HSS=19



NMME




For the rationale behind CPC outlooks see https://www.cpc.ncep.noaa.gov/products/archives/long_lead/PMD/2024/202408_PMD90D

Model Simulated/Forecast Ensemble Mean Anomalies


Model Simulated/Forecast Ensemble Average Anomalies

- AMIP simulations forced with observed sea surface temperatures (100 members ensemble)
- CFSv2 real time operational forecasts
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead). For example, 2016AMJ seasonal mean forecasts are 40 members from 22-31 March2016 initial conditions.
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013). For example, the constructed 2016AMJ seasonal mean forecasts are the average of April2016 forecasts from 22-31 March2016 initial conditions, May2016 forecasts from 21-30 April2016 initial conditions, and June2016 forecasts from 22-31 May2016 initial conditions.
- Numbers at the panels indicate the spatial anomaly correlation (AC). “w/o mean” is AC with area mean removed.


SON2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies SST(K)




SON2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)




SON2024 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)




SON2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




SON2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies Prec(mm/day)



SON2024 Observed & Model Simulated/Forecast
Ensemble Average Anomalies z200(m)
(full anomalies: shaded; eddy anomalies: contours)



SON2024 Observed & Model Simulated/Forecast Ensemble Average Anomalies T2m(K)




Model Simulated/Forecast Anomalies: Individual Runs


Model Simulated/Forecast Anomalies: Individual Runs

- In this analysis, anomalies from individual model runs are compared against the observed seasonal mean anomalies. The spatial resemblance between them is quantified based on anomaly correlation (AC).
- The distribution of AC across all model simulations is indicative of probability of observed anomalies to have a predictable (or attributable) component.
- One can also look at best and worst match between model simulated/forecast anomalies to assess the range of possible seasonal mean outcomes.
- For further details see: Kumar, A., M. Chen, M. Hoerling, and J. Eischeid (2013), Do extreme climate events require extreme forcings? *Geophys. Res. Lett.*, 40, 3440-3445. [doi:10.1002/grl.50657](https://doi.org/10.1002/grl.50657).


SON2024 Anomaly Correlation for Individual AMIP Simulation
with Observation -- $z200(20N-90N)$



Observed & AMIP Ensemble Mean Anomalies
SON2024 z200(m) 100 runs/worst 4 runs/best 4 runs
(full anomalies: shaded; eddy anomalies: contours)





SON2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- z200 (20N–90N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
SON2024 z200(m) 40 runs/worst 4 runs/best 4 runs

Seasonal Forecast

(full anomalies: shaded; eddy anomalies: contours)





Observed & CFSv2 Forecast Ensemble Average Anomalies


SON2024 z200(m) 40 runs/worst 4 runs/best 4 runs

Reconstructed Forecast


(full anomalies: shaded; eddy anomalies: contours)




SON2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- Prec(NA)/SST(30S–30N)




Observed & CFSv2 Forecast Ensemble Average Anomalies
SON2024 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
SON2024 Prec(mm/day) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast




SON2024 Anomaly Correlation for Individual CFSv2 Forecast with Observation -- T2m(NA)/SST(30S–30N)



Observed & CFSv2 Forecast Ensemble Average Anomalies
SON2024 T2m(K) 40 runs/worst 4 runs/best 4 runs
Seasonal Forecast




Observed & CFSv2 Forecast Ensemble Average Anomalies
SON2024 T2m(K) 40 runs/worst 4 runs/best 4 runs
Reconstructed Forecast



$z200(m)$ Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) SON2024 $z200(m)$ eddy & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th – 20th of the prior month.
- 4th row: 1st – 10th of the prior month.

T2m(k) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) SON2024 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th - 20th of the prior month.
- 4th row: 1st - 10th of the prior month.

Prec(mm/day) Monthly Means from Seasonal Forecast

Monthly Means from Seasonal Fcst (40ensm) SON2024 Prec(mm/day) & Obs




Top row: Observed anomaly.

CFSv2 seasonal forecasts from different initial conditions in the month prior to the target season:

- 2nd row: last 10 days of the prior month.
- 3rd row: 11th – 20th of the prior month.
- 4th row: 1st – 10th of the prior month.

$z200(m)$ Monthly Means from Monthly Forecast




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

T2m(k) Monthly Means from Monthly Forecast

Monthly Means from Monthly Fcst SON2024 T2m(K) & Obs




Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.

Prec(/mm/day) Monthly Means from Monthly Forecast



Top row: Observed anomaly.

CFSv2 monthly forecasts from different initial conditions in the month prior to the target month:

- 2nd row: last 3 days of the prior month.
- 3rd row: 18th – 20th of the prior month.
- 4th row: 7th – 10th of the prior month.



Seasonal Forecasts from Multi-Model Ensemble Systems

- WMO Lead Center for Long-Range Forecast Multi-Model Ensemble (LC-LRFMME).
<https://www.wmclc.org/>
- Copernicus Climate Change Service (C3S) Multi-model seasonal forecasts.
https://climate.copernicus.eu/charts/c3s_seasonal/
- North American Multi-Model Ensemble (NMME) seasonal forecasts.
<https://www.cpc.ncep.noaa.gov/products/NMME/>



LC-LRFMM Seasonal Forecasts

(<https://www.wmolk.org/>)

Ensemble means




Probabilities



C3S Seasonal Forecast

(https://climate.copernicus.eu/charts/c3s_seasonal/)



PROGRAMME OF
THE EUROPEAN UNION



IMPLEMENTED BY



PROGRAMME OF
THE EUROPEAN UNION




IMPLEMENTED BY




North American Multi-Model Ensemble Seasonal Forecast

(<https://www.cpc.ncep.noaa.gov/products/NMME/>)






200mb Height from Linear Model

SON2024 200mb Eddy HGT(m)
OBS vs. Linear Model Response to Tropical Heating
Heating is converted from Prate in 15S–15N



Seasonal Forecasts from the Constructed Analog Model



Background & Methodology

Attribution of Seasonal Climate Anomalies

- Goal
 - In the context of prediction of seasonal climate variability, utilize seasonal climate forecasts and atmospheric general circulation model (AGCM) simulations to attribute possible causes for the observed seasonal climate anomalies.
 - The analysis can also be considered as an analysis of predictability of the observed seasonal climate anomalies.

Methodology - 1

- Compare observed seasonal mean anomalies with those from model simulations and forecasts.
- Ensemble averaged model simulated/predicted seasonal mean anomalies are an indication of the predictable (or attributable) component of the corresponding observed anomalies.
- For seasonal mean atmospheric anomalies, predictability could be due to
 - Anomalous boundary forcings [e.g., sea surface temperature (SSTs); soil moisture etc.];
 - Atmospheric initial conditions.
- The influence of anomalous boundary forcings (particularly due to SSTs, can be inferred from the ensemble mean of AGCM simulations forced by observed SSTs, the so called AMIP simulations). This component of predictability (or attributability) is more relevant for longer lead seasonal forecasts.

Methodology - 2

- The influence of the atmospheric initial state can be inferred from initialized predictions. This component is more relevant for short lead seasonal forecasts.
- The influence of unpredictable component in the atmospheric variability can be assessed from the analysis of individual model simulations, and the extent anomalies in individual runs deviate from the ensemble mean anomalies.
- The relative amplitude of ensemble averaged seasonal mean anomalies to the deviations of seasonal mean anomalies in the individual model runs from the ensemble average is a measure of seasonal predictability (or the extent observed anomalies are attributable).
- Observed anomalies are equivalent to a realization of a single model run, and therefore, analysis of individual model runs also gives an appreciation of how much observed anomalies can deviate from the component that is attributable (Kumar et al. 2013).

Data

- Observations
 - SST: OI version 2 analysis (Reynolds et al., 2007)
 - Prec: CMAP monthly analysis (Xie and Arkin, 1997)
 - T2m: GHCN-CAMS land surface temperature monthly analysis (Fan and van den Dool, 2008)
 - 200mb height (z200): CFSR (Saha et al., 2010)
- 0-month-lead seasonal mean forecasts from CFSv2 (Saha et al. 2014)
 - Seasonal forecast: the seasonal mean forecasts based on 40 members from the latest 10 days before the target season (0-month-lead);
 - Reconstructed forecast: the seasonal mean forecasts constructed from 3 individual monthly forecasts with the latest 10 days initial conditions for each individual monthly forecasts. This approach for constructing seasonal mean anomalies has more influence from the initial conditions (Kumar et al. 2013);
- Seasonal mean AMIP simulation based on GFS_FV3 (provided by Dr. Tao Zhang/CPC)
 - 100 members
- All above seasonal mean anomalies are based on 1991-2020 climatology.
- z200 responses to tropical heating in linear model.
- Seasonal mean anomalies of z200, T2m, and Prec forecasted from the Constructed Analog Model.