



Madden-Julian Oscillation: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
June 22, 2015**



Outline

- **Overview**
- **Recent Evolution and Current Conditions**
- **MJO Index Information**
- **MJO Index Forecasts**
- **MJO Composites**



Overview

- **The MJO remains active with the enhanced convective phase currently centered over the Maritime Continent.**
- **Destructive interference between the MJO and the El Niño state continued over the past week with the enhanced phase of the MJO currently disrupting El Niño-favored suppressed convection over the Maritime continent. Enhanced convection has returned to the central Pacific.**
- **Dynamical model MJO index forecasts favor continued eastward propagation of the MJO at varying amplitudes to the western Pacific during the upcoming two weeks.**
- **The MJO is expected to play a substantial role in the global tropical convective pattern during the period and favors an enhancement of the South Asian Monsoon while suppressed convection is more likely along the equatorial Indian Ocean and some parts of Central America, Mexico and the Caribbean. Tropical cyclone activity is favored for portions of the western Pacific.**

Additional potential impacts across the global tropics and a discussion for the U.S. are available at:
<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/ghazards/index.php>




850-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies developed over southern Asia and the far western Pacific during the last five days.

Winds have been close to average over the later five days across the central Pacific.

Westerly anomalies strengthened over the eastern tropical Atlantic.




850-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

CDAS 850-hPa U Anoms. (5N-5S)

Time



During December, easterly anomalies persisted from 120E to the Date Line (red box).

Westerly anomalies associated with the MJO propagated eastward (dashed line) during the first half of January. Westerly anomalies returned to the Western Pacific during late January and early February.

Strong westerly anomalies associated with the MJO, an equatorial Rossby wave (ERW) and El Niño base state conditions resulted in strong westerly anomalies propagating west of the Date Line during early March.


During April and May, westerly anomalies expanded over much of the central and eastern Pacific, consistent with El Niño. More recently, eastward propagation on the intraseasonal time scale was observed.

Longitude



OLR Anomalies – Past 30 days

Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)
Wetter-than-normal conditions, negative OLR anomalies (blue shading)




Convection weakened near the Date Line during late May, with enhanced convection persisting over the eastern Pacific. Suppressed convection continued over southeastern Asia, the Maritime Continent, and the western Pacific.

During early June, enhanced convection over the eastern Pacific and Arabian Sea was in part associated with tropical cyclone activity. The low frequency suppressed convection signal weakened over the southern Maritime Continent, but persisted over Southeast Asia.

By mid-June, enhanced convection returned to the western and central Pacific. Enhanced convection was also evident from India across the Maritime continent to the western Pacific due to the MJO.



Outgoing Longwave Radiation (OLR) Anomalies (2.5°N-17.5°N)



Drier-than-normal conditions, positive OLR anomalies (yellow/red shading)

Wetter-than-normal conditions, negative OLR anomalies (blue shading)

(Courtesy of CAWCR Australia Bureau of Meteorology)

The MJO became active and strong during March, with eastward propagation of enhanced (suppressed) convective anomalies evident across the Pacific (Indian Ocean and Maritime Continent).

Since late March, enhanced (suppressed) convection has dominated at or east of the Date Line (Maritime Continent) (red box), consistent with El Niño conditions. Kelvin Waves were the most prominent subseasonal features during April and May.

During late May and June, slower, more robust eastward propagation was evident, consistent with MJO activity.




200-hPa Velocity Potential Anomalies (5°S-5°N)

Positive anomalies (brown shading) indicate unfavorable conditions for precipitation

Negative anomalies (green shading) indicate favorable conditions for precipitation

Time
↓



Longitude

From December 2014 though January 2015, the MJO was active, as indicated by eastward propagation of alternating anomalies. At times, the signal was dominated by faster-moving variability (likely Kelvin Wave activity).

The signal was weak much of January and during February.

During March, MJO activity was observed, with anomalies becoming strong as they interacted with the developing low frequency state.

Negative anomalies persisted near the Date Line and to the east since early April due to the El Niño base state. During this time, Kelvin wave activity (fast eastward propagation) has been the primary subseasonal mode of variability evident in this field.


More recently, slower eastward propagation of positive VP anomalies was observed over the West Pacific, while negative anomalies developed over the Indian Ocean, consistent with MJO activity.



IR Temperatures (K) / 200-hPa Velocity Potential Anomalies

Positive anomalies (brown contours) indicate unfavorable conditions for precipitation

Negative anomalies (green contours) indicate favorable conditions for precipitation



The spatial velocity potential pattern has a coherent Wave-1 structure, with positive anomalies (associated with suppressed convection) over the Atlantic and Africa with negative anomalies (associated with enhanced convection) over the eastern Indian Ocean and Maritime Continent. Enhanced convection over the Pacific is associated with a Kelvin Wave emerging ahead of the primary MJO envelope constructively interacting with the base state.




200-hPa Vector Wind Anomalies (m s^{-1})

Note that shading denotes the zonal wind anomaly

Blue shades: Easterly anomalies

Red shades: Westerly anomalies



Westerly anomalies have strengthened over the Americas during the last five days (black box).




200-hPa Zonal Wind Anomalies (m s^{-1})

Westerly anomalies (orange/red shading) represent anomalous west-to-east flow

Easterly anomalies (blue shading) represent anomalous east-to-west flow

Time
↓



Longitude

During late December through the mid-April, westerly anomalies increased in coverage and intensity from 120W to 80W.


Westward propagation of westerly anomalies was evident over the eastern Pacific during late February and again in March (black arrows).

Easterly anomalies have generally persisted over the central and eastern Pacific (red box) consistent with El Nino since early May.

More recently, westerly anomalies propagated eastward from the Maritime Continent to the western Hemisphere, consistent with MJO activity.



Weekly Heat Content Evolution in the Equatorial Pacific



Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.

The upwelling phase of a Kelvin wave went through during May-July 2014.

During October-November, positive subsurface temperature anomalies increased and shifted eastward in association with the downwelling phase of a Kelvin wave.

During November - January, the upwelling phase of a Kelvin wave shifted eastward.

During January through April, a very strong downwelling Kelvin Wave was observed..

Positive anomalies persisted over the central and Eastern Pacific, with evidence of a potential second downwelling Kelvin Wave evident during late May and early June.



MJO Index -- Information

- The MJO index illustrated on the next several slides is the CPC version of the Wheeler and Hendon index (2004, hereafter WH2004).

Wheeler M. and H. Hendon, 2004: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Monthly Weather Review*, 132, 1917-1932.


- The methodology is very similar to that described in WH2004 but does not include the linear removal of ENSO variability associated with a sea surface temperature index. The methodology is consistent with that outlined by the U.S. CLIVAR MJO Working Group.

Gottschalck et al. 2010: A Framework for Assessing Operational Madden-Julian Oscillation Forecasts: A CLIVAR MJO Working Group Project, *Bull. Amer. Met. Soc.*, 91, 1247-1258.

- The index is based on a combined Empirical Orthogonal Function (EOF) analysis using fields of near-equatorially-averaged 850-hPa and 200-hPa zonal wind and outgoing longwave radiation (OLR).



MJO Index -- Recent Evolution




- The axes (RMM1 and RMM2) represent daily values of the principal components from the two leading modes
- The triangular areas indicate the location of the enhanced phase of the MJO
- Counter-clockwise motion is indicative of eastward propagation. Large dot most recent observation.
- Distance from the origin is proportional to MJO strength
- Line colors distinguish different months

Eastward propagation of a MJO signal was evident during the past two weeks, with the enhanced phase centered over the Maritime continent.



MJO Index – Historical Daily Time Series



Time series of daily MJO index amplitude from 2007 to present.

Plot puts current MJO activity in recent historical context.




Ensemble GFS (GEFS) MJO Forecast

Yellow Lines – 20 Individual Members
Green Line – Ensemble Mean

RMM1 and RMM2 values for the most recent 40 days and forecasts from the ensemble Global Forecast System (GEFS) for the next 15 days

light gray shading: 90% of forecasts
dark gray shading: 50% of forecasts

The GFS ensemble MJO index forecast depicts continued slow eastward propagation of a MJO signal during the next two weeks to the western Pacific.







Ensemble Mean GFS MJO Forecast

Figures below show MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days




The ensemble GFS forecast depicts enhanced convection slowly shifting eastward from southern Asia and the Maritime continent to the western and central Pacific over the next two weeks. Suppressed convection is forecast for portions of Central America and later the Indian Ocean over the period.




Constructed Analog (CA) MJO Forecast

Figure below shows MJO associated OLR anomalies only (reconstructed from RMM1 and RMM2) and do not include contributions from other modes (*i.e.*, ENSO, monsoons, etc.)

Spatial map of OLR anomalies for the next 15 days



Time-longitude section of (7.5°S-7.5°N) OLR anomalies for the last 180 days and for the next 15 days





The statistical forecast depicts a similar, but weaker MJO signal strength as compared to the ensemble GFS over the two week period.



MJO Composites – Global Tropics

850-hPa Velocity Potential and
Wind Anomalies (May-Sep)

Precipitation Anomalies (May-Sep)






U.S. MJO Composites – Temperature

Left hand side plots show temperature anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Blue (orange) shades show negative (positive) anomalies respectively.

Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.




Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>



U.S. MJO Composites – Precipitation

- Left hand side plots show precipitation anomalies by MJO phase for MJO events that have occurred over the three month period in the historical record. Brown (green) shades show negative (positive) anomalies respectively.
- Right hand side plots show a measure of significance for the left hand side anomalies. Purple shades indicate areas in which the anomalies are significant at the 95% or better confidence level.



Zhou et al. (2011): A composite study of the MJO influence on the surface air temperature and precipitation over the Continental United States, *Climate Dynamics*, 1-13, doi: 10.1007/s00382-011-1001-9

<http://www.cpc.ncep.noaa.gov/products/precip/CWlink/MJO/mjo.shtml>